Bull. Aust. Math. Soc. **106** (2022), 132–143 doi:10.1017/S0004972722000144

LI-YORKE CHAOS ALMOST EVERYWHERE: ON THE PERVASIVENESS OF DISJOINT EXTREMALLY SCRAMBLED SETS

LIUCHUN DENG[®], M. ALI KHAN[®] and ASHVIN V. RAJAN[®]

(Received 22 September 2021; accepted 30 December 2021; first published online 3 March 2022)

Abstract

We show that there exists a continuous function from the unit Lebesgue interval to itself such that for any $\epsilon \ge 0$ and any natural number *k*, any point in its domain has an ϵ -neighbourhood which, when feasible, contains *k* mutually disjoint extremally scrambled sets of identical Lebesgue measure, homeomorphic to each other. This result enables a satisfying generalisation of Li–Yorke (topological) chaos and suggests an open (difficult) problem as to whether the result is valid for piecewise linear functions.

2020 Mathematics subject classification: primary 37E05; secondary 37N40, 91B55, 92D25.

Keywords and phrases: scrambled set, extremally scrambled set, Smith–Volterra–Cantor set, Lebesgue measure, Li–Yorke chaos, topological chaos.

1. Introduction

It was elementary, but not simple by any means. The problem turned out to be deep, the appearance of simplicity was deceptive. Khinchin [8]

Independent and pioneering results of Sharkovsky [17] and Li and Yorke [9] have led to a rich literature on continuous self-maps on the unit interval, and the phrase 'Li–Yorke chaotic maps' is now part of the vernacular of dynamical systems and affiliated applied fields, not only physics and engineering, but also evolutionary biology and economics. Sharkovsky defines an order relation on the possible periods of periodic points, establishing precise conditions under which such interval maps can have points of any period, while Li and Yorke furnish conditions under which nonperiodic chaotic behaviour is displayed by an uncountable subset of initial points of an interval map. Much effort has been expended to capture the apparently random behaviour of chaotic trajectories through precise mathematical definitions of chaos itself. The excellent monograph by Ruette [16] contains a very precise taxonomy of the different notions of chaos that have been explored (see page VIII of the introduction for a clear schematic



Liuchun Deng acknowledges the support of a start-up grant from Yale-NUS College.

[©] The Author(s), 2022. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.

summary). For the distinctions between turning points, periodic points, periodic orbits, orbit portraits and their evolution under the general rubric of the concept of an *orbit*, see Milnor and Thurston's monograph [10] and the extensive work that has followed it: the field of economic dynamics still has much to learn from this literature.

The result we present belongs more generally to the subject of the pervasiveness of Li–Yorke chaos, the existence of scrambled sets in the domain of a map that can be arbitrarily large or arbitrarily small as measured by the Lebesgue measure but are pervasive in being spread all over the domain. The result addresses the 'observability' of chaos exhibited by interval maps as tracked by the Lebesgue measure of sets of initial points with chaotic trajectories (see Misiurewicz [11] for the linkage between the Lebesgue measure of scrambled sets and observability of chaos). As background, let us note Jiménéz-López singling out in his review [5] of Ruette [16] 'a very beautiful (and far from easy to prove) dichotomy result by Smítal: a continuous interval map is either Li–Yorke chaotic, or all points are (up to small errors) eventually periodic for it'.

In this connection, Nathanson [13] presents a continuous self-map on the interval that exhibits unobservable Li–Yorke chaos in a system exhibiting dynamics that can be made as regular (periodic) as one pleases. For any natural number $n \ge 3$, he constructs a piecewise linear continuous map such that all trajectories except for a set of Lebesgue measure zero have eventually periodic trajectories of period n. On the other hand, there is a dual, apparently orthogonal, result due to Kan [6], which questions the prevailing sentiment of the negligibility of the set of initial conditions that give rise to Li–Yorke chaotic trajectories. His canonical example consists of a continuous self-map defined on the unit interval which has a scrambled set whose Lebesgue measure is 1/8. This result was soon generalised to show a scrambled set, first of full outer Lebesgue measure [18] and then [12] and [15] for extensions and [1], [2, footnote 66], [14, page 63] and [16, page 111] for its framing more broadly in the current literature.

The point of this paper (the question posed and answered) is that we can extend Kan's construction to find a chaotic interval map that has an arbitrarily large (but finite) number of mutually disjoint extremally scrambled sets which are homeomorphic to each other, and each of which has positive Lebesgue measure. *Extremally scrambled* sets of functions defined on the closed unit interval are scrambled sets in the Li–Yorke sense, which are comprised of initial points whose trajectories contain subsequences whose distance from an arbitrary periodic point can be bounded below by 1/2 (the largest possible positive constant in this context). A precise formal definition is given in Section 2. Our first theorem shows that extremally scrambled sets are dispersed all over the domain of a continuous self-map of the interval. Our second theorem, based on a similar construction, allows us to recover not only Kan's result but also the theorems of Smítal and Rajan as particular cases. Equally to the point, our construction yields, with straightforward modifications, a continuous self-map whose scrambled set, though again dispersed all over the interval, can be of zero Lebesgue measure.

In terms of applications, and staying with economic dynamics, let us note that Nathanson's map with parallel left and right arms was extended, by Khan and

[3]

Rajan [7], to yield a family of maps whose arms are not parallel, and thereby shown to constitute reduced forms of workhorse two-sector models in mathematical economics. In [3], the application is shown to go beyond the technical into the substantive. The Khan–Rajan extension of Nathanson shows how in equilibrium growth 'anything goes' in the sense that, for any natural number n, there exists a prototypical two-sector model, associated with Shinkai, Uzawa and Srinivasan, that, while admitting Li–Yorke chaotic trajectories, exhibits eventually periodic trajectories of period n, starting from almost any initial condition. Equally importantly, while this is evident in (Keynesian) models of equilibrium growth, such trajectories are rendered impossible in (Ramseyan) models of optimal growth in which intertemporal arbitrage conditions play a determining role (see [3]).

Before turning to the plan of the work and concluding this introduction, let us note some questions that are rendered open by our result: the open questions are arguably as important as the answers given here. The first is the observation that the map we construct, like that of [6], barring the two linear arms on left and right outside, has Cantor's 'devil's staircase' in the middle. We can then ask whether the parallelism of the left and right arms, reminiscent of Nathanson's map, can be relaxed to nonparallel arms, \dot{a} la Khan and Rajan [7], to strengthen the results on the two-sector model available in [3]. Technically, this would also strengthen the result of [15]. However, as stated above, this is not merely of technical interest, but of substantive consequence for economic dynamics.

The second open question relates to an orthogonal direction of unobservable Li–Yorke chaos. Can Corollary 2.7 below be extended to show that for any natural numbers $n, m \in \mathbb{N}$, and for any piecewise linear map with the number of pieces less than or equal to *n*, the size of the *m* dispersed scrambled sets is zero?

Finally, note that the our work has nothing to say about the closely related result of Misiurewicz [12] which uses topological conjugacy, and thereby different techniques, to obtain a continuous map for which there exists a scrambled set of Lebesgue measure 1. Progress in this direction would lead to a fuller exploration in economic dynamics of an idea that has not been sufficiently attended to on the topological conjugacy of economic models, and thereby an invariance of the qualitative properties of their dynamics.

After this introduction, Section 2 presents the results and Section 3 the proofs. Section 3 is divided into three subsections: the argument, the construction and the proofs of the ancillary results.

2. The result

Denote by C([0, 1]) the space of continuous functions from the unit interval to itself.

DEFINITION 2.1. An uncountable subset $K \subset [0, 1]$ is called an *extremally scrambled* set for a function $f \in C([0, 1])$, if, for any $x, y \in K, x \neq y$:

Disjoint extremally scrambled sets

(i) $\limsup_{k \to \infty} |f^k(x) - f^k(y)| = 1;$

- (ii) $\lim \inf_{k \to \infty} |f^k(x) f^k(y)| = 0$; and
- (iii) $\limsup_{k\to\infty} |f^k(x) f^k(x_0)| \ge \frac{1}{2}$ for any periodic point x_0 of f.

THEOREM 2.2. For any natural number k and any positive real number ϵ , there exists a continuous mapping $f_{k,\epsilon} \in C([0, 1])$ such that for any $x \in [0, 1]$, the interval $(x - \epsilon, x + \epsilon) \cap [0, 1]$ contains k mutually disjoint extremally scrambled sets of identical positive Lebesgue measure, homeomorphic to each other.

If we set k = 1 in Theorem 2.2, following a similar construction to Section 3.2 below, we can adjust the size of the extremally scrambled set to obtain the following result.

THEOREM 2.3. For any $(\epsilon, \delta) \in (0, 1) \times (0, 1]$, there exists a continuous mapping $g_{\epsilon,\delta} \in C([0, 1])$ that has an extremally scrambled set of Lebesgue measure $(1 - \delta)$, such that for any $x \in [0, 1]$, the interval $(x - \epsilon, x + \epsilon) \cap [0, 1]$ contains an extremally scrambled set having positive Lebesgue measure.

On letting the Lebesgue measure of the union of the extremally scrambled sets be any real number in [0, 1), Theorem 2.3 can be sharpened to yield the results in Kan [6], Smítal [18] and Rajan [15] as corollaries. It should be noted that in contrast to the proofs in [6, 15] which influence the methods used herein, Smítal obtains his result through a very different construction.

COROLLARY 2.4 (Kan). There exists $f \in C([0, 1])$ which possesses an extremally scrambled set of Lebesgue measure 1/8.

COROLLARY 2.5 (Smítal). For any $\epsilon \in (0, 1)$, there exists $f_{\epsilon} \in C([0, 1])$ which possesses a scrambled set of Lebesgue measure greater than $1 - \epsilon$.

COROLLARY 2.6 (Rajan). For any $\epsilon \in (0, 1)$, there exists $f_{\epsilon} \in C([0, 1])$ which possesses an extremally scrambled set of Lebesgue measure $1 - \epsilon$.

COROLLARY 2.7. For any $\epsilon \in (0, 1)$, there exists a mapping $f_{\epsilon} \in C([0, 1])$ that has an extremally scrambled set of Lebesgue measure 0, such that for any $x \in [0, 1]$, the interval $(x - \epsilon, x + \epsilon) \cap [0, 1]$ contains an extremally scrambled set having Lebesgue measure 0.

3. The proofs

The proof of Theorem 2.2 rests on Proposition 3.1 stated and proved below. The proof of Theorem 2.3 follows similarly from a straightforward modification of Proposition 3.1 in the light of Remarks 3.2 and 3.3 in Section 3.2 below. The proofs of the four corollaries are straightforward and left to the reader.

[4]

3.1. The proof of Theorem 2.2. For any arbitrary positive integer *m*, define

$$\mathcal{L}_{m} = \begin{bmatrix} 0, \frac{1}{3^{m}} \end{bmatrix}, \quad \mathcal{M}_{m} = \begin{bmatrix} \frac{1}{3^{m}}, 1 - \frac{1}{3^{m}} \end{bmatrix}, \quad \mathcal{R}_{m} = \begin{bmatrix} 1 - \frac{1}{3^{m}}, 1 \end{bmatrix}, \quad \mathcal{M}_{m}^{\circ} = \left(\frac{1}{3^{m}}, 1 - \frac{1}{3^{m}}\right),$$
$$\mathcal{G}_{m} = \left\{ g_{m} \in \mathcal{C}([0, 1]) : g_{m}(x) = \begin{cases} 3^{m}x & \text{if } x \in \mathcal{L}_{m} \\ 3^{m}x - (3^{m} - 1) & \text{if } x \in \mathcal{R}_{m} \end{cases} \right\}.$$

For any arbitrary positive integer *n*, partition \mathcal{M}_m° into 2^n mutually disjoint intervals of length $\delta_m^n = 2^{-n} \mu(\mathcal{M}_m^\circ)$ with each interval defined as

$$M_{m,j}^{n} = \left(\frac{1}{3^{m}} + (j-1)\delta_{m}^{n}, \frac{1}{3^{m}} + j\delta_{m}^{n}\right), \text{ for } 1 \le j \le 2^{n}.$$

PROPOSITION 3.1. For any positive integers m, n and k, there exists a continuous mapping in \mathcal{G}_m such that each interval $M_{m,j}^n$ (for $1 \le j \le 2^n$) contains k mutually disjoint extremally scrambled sets of identical positive Lebesgue measure, homeomorphic to each other.

PROOF OF THEOREM 2.2. Choose positive integers *m* and *n* sufficiently large so that $3^{-m} + \delta_m^n < \epsilon$. From Proposition 3.1, there exists a continuous mapping *f* in \mathcal{G}_m such that each interval $M_{m,j}^n$ (for $1 \le j \le 2^n$) contains *k* mutually disjoint extremally scrambled sets of identical positive Lebesgue measure, homeomorphic to each other. Define $f_{k,\epsilon} = f$. Then, $[0, \epsilon)$ contains $M_{m,1}^n$ and $(1 - \epsilon, 1]$ contains $M_{m,2^n}^n$. For any $x \in (0, 1)$, the segment $(x - \epsilon, x + \epsilon) \cap [0, 1]$ contains some $M_{m,j}^n$ and hence contains *k* mutually disjoint extremally scrambled sets of identical positive Lebesgue measure, homeomorphic to each other.

3.2. The construction. The construction is obtained through the following three steps. In step 1, divide the unit interval into three exhaustive segments \mathcal{L}_m , \mathcal{M}_m and \mathcal{R}_m as defined above. We first construct 2^n mutually disjoint sets in \mathcal{L}_m which are extremally scrambled sets for any function in \mathcal{G}_m . Then, in step 2, for the middle interval \mathcal{M}_m , we construct 2^n mutually disjoint Smith–Volterra–Cantor (SVC) sets each of which is contained in each of the 2^n intervals $\mathcal{M}_{m,j}^n$ ($1 \le j \le 2^n$). Finally, in step 3, we construct a homeomorphism from the 2^n SVC sets to the 2^n extremally scrambled sets. Extending the homeomorphism to the unit interval gives the desired continuous map $f_{m,n}$.

Step 1. Denote by Ω the space of infinite binary sequences, which is an ordered topological space when equipped with the lexicographic ordering and topologised with the topology of pointwise convergence. Consider the 2^{n+1} binary (n + 1)-tuples

$$\varepsilon_j = (e_1^j, \dots, e_{n+1}^j) \text{ for } 1 \le j \le 2^{n+1}, \text{ where } e_k^j \in \{0, 1\}.$$

Define a binary relation 0 < 1 and order the 2^{n+1} binary (n + 1)-tuples lexicographically so that $\varepsilon_1 < \varepsilon_2 < \cdots < \varepsilon_{2^{n+1}}$. Define

$$\Omega_{\varepsilon_i} = \{(\omega_k)_{k\geq 1} \in \Omega : (\omega_k)_{k\geq 1} = (\varepsilon_j, \omega_{n+2}, \omega_{n+3}, \ldots)\}.$$

137

Then $\Omega_{\varepsilon_1} < \cdots < \Omega_{\varepsilon_{2^{n+1}}}, \Omega_{\varepsilon_i} \cap \Omega_{\varepsilon_j} = \emptyset$ for $i \neq j$ and $\Omega = \Omega_{\varepsilon_1} \cup \cdots \cup \Omega_{\varepsilon_{2^{n+1}}}$. Hence the subsets $\{\Omega_{\varepsilon_1}, \ldots, \Omega_{\varepsilon_{2^{n+1}}}\}$ form a partition of the space Ω into 2^{n+1} mutually disjoint subsets. Consider the shift mapping $\sigma : \Omega \to \Omega$ given by

$$\sigma(\omega) = (\sigma(\omega_1), \sigma(\omega_2), \ldots), \text{ where } \sigma(\omega_i) = \omega_{i+1} \text{ for } j \ge 1 \text{ and } \omega \in \Omega.$$

Define $\phi_m : \Omega \to [0, 1]$ by

$$\phi_m(\omega) = \sum_{k=1}^{\infty} \frac{(3^m - 1)\omega_k}{3^{mk}},$$

and observe the following routine facts:

- (i) ϕ_m is an order-preserving homeomorphism;
- (ii) $\phi_m(\Omega) \cap \mathcal{M}_m^\circ = \emptyset$; and
- (iii) $\phi_m(\Omega)$ is homeomorphic to the classical 'middle-thirds' Cantor set [4].

Consider $(\omega_k)_{k\geq 1} \in \Omega_{\varepsilon_i}$ for $1 \leq j \leq 2^n$. For any positive integer *t*, define

$$\Lambda_t = \left(\underbrace{\omega_1 \cdots \omega_1}_{t \text{ times}} \underbrace{\omega_2 \cdots \omega_2}_{t \text{ times}} \cdots \underbrace{\omega_t \cdots \omega_t}_{t \text{ times}}\right) \text{ and } B_t = \left(\underbrace{0 \cdots 0}_{t \text{ times}} \underbrace{1 \cdots 1}_{t \text{ times}}\right).$$

The mapping $Z_{\varepsilon_i} : \Omega_{\varepsilon_i} \to Z_{\varepsilon_i}(\Omega_{\varepsilon_i}) \subset \Omega_{\varepsilon_i}$ defined by

$$Z_{\varepsilon_j}(\varepsilon_j,\omega_{n+2},\ldots)=(\varepsilon_j,B_1\Lambda_1B_2\Lambda_2B_3\Lambda_3\cdots)$$

is an order-preserving homeomorphism. We define

$$K_{\varepsilon_j} = \phi_m \circ Z_{\varepsilon_j}(\Omega_{\varepsilon_j}), \text{ for } 1 \le j \le 2^n,$$

and observe that as $\Omega_{\varepsilon_1} < \cdots < \Omega_{\varepsilon_{2n}}$ and $Z_{\varepsilon_j}(\Omega_{\varepsilon_j}) \subset \Omega_{\varepsilon_j}$ for $1 \le j \le 2^n$, we have $Z_{\varepsilon_1}(\Omega_{\varepsilon_1}) < \cdots < Z_{\varepsilon_{2n}}(\Omega_{\varepsilon_{2n}})$. Since the function ϕ_m is an order-preserving homeomorphism and since $\phi_m(\Omega_{\varepsilon_j}) \subset \mathcal{L}_m$ for $1 \le j \le 2^n$, it follows that $\phi_m(Z_{\varepsilon_1}(\Omega_{\varepsilon_1})) = K_{\varepsilon_1} < \cdots < \phi_m(Z_{\varepsilon_{2n}}(\Omega_{\varepsilon_{2n}})) = K_{\varepsilon_{2n}}$. Thus, the sets $K_{\varepsilon_1}, \ldots, K_{\varepsilon_{2n}}$ are a collection of 2^n mutually disjoint subsets of the interval \mathcal{L}_m . Later we will establish in Proposition 3.6 that each of the 2^n mutually disjoint subsets K_{ε_j} is an extremally scrambled set for every function $g_m \in \mathcal{G}_m$.

Step 2. For ease of exposition, let $M_{\varepsilon_j} = (\alpha_j, \beta_j) \equiv M_{m,j-2^n}^n$ for $2^n + 1 \le j \le 2^{n+1}$. We construct positive measure SVC sets $C_{\varepsilon_j} \subset M_{\varepsilon_j}$ such that $\mu(C_{\varepsilon_j}) = 1/2^{n+2}$. To this end, consider any one of the above intervals M_{ε_j} , and consider its midpoint $(\alpha_j + \beta_j)/2 = m_j$. Choose a positive real number ϵ such that

$$\frac{1}{2^{n+2}} < \epsilon < \delta_m^n$$
 and let $\delta = \left(\epsilon - \frac{1}{2^{n+2}}\right)$

Define $E_{\varepsilon_j}^0 = [\ell_{\varepsilon_j}, r_{\varepsilon_j}] = [m_j - \frac{1}{2}\epsilon, m_j + \frac{1}{2}\epsilon]$ and set $\ell_{\varepsilon_j0} = \ell_{\varepsilon_j} + \frac{1}{4}\delta, r_{\varepsilon_j0} = m_j, \ell_{\varepsilon_j1} = m_j + \frac{1}{4}\delta, r_{\varepsilon_j1} = r_{\varepsilon_j}$ and $E_{\varepsilon_j}^1 = [\ell_{\varepsilon_j0}, r_{\varepsilon_j0}] \cup [\ell_{\varepsilon_j1}, r_{\varepsilon_j1}]$. Observe that $\mu(E_{\varepsilon_j}^1) = \mu(E_{\varepsilon_j}^0) - \frac{1}{2}\delta$. Next, set $\ell_{\varepsilon_j00} = \ell_{\varepsilon_j0} + \frac{1}{16}\delta, r_{\varepsilon_j00} = \frac{1}{2}(\ell_{\varepsilon_j0} + r_{\varepsilon_j0}), \ell_{\varepsilon_j01} = r_{\varepsilon_j00} + \frac{1}{16}\delta, r_{\varepsilon_j01} = r_{\varepsilon_j0}, \ell_{\varepsilon_j10} = \ell_{\varepsilon_j1} + \frac{1}{16}\delta, r_{\varepsilon_j10} = \frac{1}{2}(\ell_{\varepsilon_j1} + r_{\varepsilon_j1}), \ell_{\varepsilon_j11} = r_{\varepsilon_j10} + \frac{1}{16}\delta, r_{\varepsilon_j11} = r_{\varepsilon_j1}$ and define $E_{\varepsilon_j}^2 = [\ell_{\varepsilon_j00}, r_{\varepsilon_j00}] \cup$

 $[\ell_{\varepsilon_j01}, r_{\varepsilon_j01}] \cup [\ell_{\varepsilon_j10}, r_{\varepsilon_j10}] \cup [\ell_{\varepsilon_j11}, r_{\varepsilon_j11}].$ Observe that $\mu(E_{\varepsilon_j}^2) = \mu(E_{\varepsilon_j}^1) - \frac{1}{2}\delta = \mu(E_{\varepsilon_j}^0) - (\frac{1}{2}\delta + \frac{1}{4}\delta).$ Repeating this construction yields a decreasing sequence of sets $E_{\varepsilon_j}^k$ for k = 0, 1, 2, ... in which every $E_{\varepsilon_j}^k$ is a finite union of disjoint closed and bounded intervals and is therefore compact. Observe that all the right end points of $E_{\varepsilon_j}^k$ are retained in $E_{\varepsilon_j}^{k+1}$, but that all the left end points of $E_{\varepsilon_j}^k$ have been replaced by new left end points which only survive one iteration of this construction of the SVC set. Thus, the decreasing sequence of nonempty compact sets $E_{\varepsilon_j}^0 \supset E_{\varepsilon_j}^1 \supset E_{\varepsilon_j}^2 \supset \cdots$ must have a nonempty intersection $C_{\varepsilon_j} = \bigcap_{k=0}^{\infty} E_{\varepsilon_j}^k \neq \emptyset$ where

$$\mu(C_{\varepsilon_j}) = \mu(E^0_{\varepsilon_j}) - \sum_{k=1}^{\infty} \frac{\delta}{2^k} = \mu(E^0_{\varepsilon_j}) - \delta = \epsilon - \delta = \epsilon - \left(\epsilon - \frac{1}{2^{n+2}}\right) = \frac{1}{2^{n+2}}$$

Since $C_{\varepsilon_j} \subset M_{\varepsilon_j}$, the 2^n mutually disjoint SVC sets $C_{\varepsilon_{2^{n+1}}}, \ldots, C_{\varepsilon_{2^{n+1}}}$ in \mathcal{M}_m° all have positive Lebesgue measure $1/2^{n+2}$.

REMARK 3.2. For any $\delta \in (0, 1)$, pick *m* such that the length of the middle interval is strictly greater than $1 - \delta$. We can adjust the measure of what we remove from each interval M_{ε_j} to make sure that the sum of the measures of the SVC sets adds up to $1 - \delta$.

REMARK 3.3. We can also make each Cantor set in our construction have measure 0 by the standard procedure of iteratively removing the open middle third from a set of intervals. This construction will yield pervasive scrambled sets of measure 0 as stated in Corollary 2.7.

Step 3. Fix any particular j with $2^n + 1 \le j \le 2^{n+1}$. We shall construct an order-preserving homeomorphism $\xi_{\varepsilon_j} : \Omega_{\varepsilon_j} \to C_{\varepsilon_j}$, where Ω_{ε_j} is constructed in step 1 and C_{ε_j} is constructed in step 2. Towards this end, for every positive integer k > 1 and for any (k - 1)-tuple $(\omega_1, \ldots, \omega_{k-1})$, replace in Ω

 $(\omega_1,\ldots,\omega_{k-1},1,0,0,\ldots,0,0,\ldots)$ by $(\omega_1,\ldots,\omega_{k-1},0,1,1,\ldots,1,1,\ldots),$

where the first terminates in an infinite sequence of 0s and the second terminates in an infinite sequence of 1s.

Consider the set $\Omega_{\varepsilon_j}^1$ of all binary sequences in Ω_{ε_j} that terminate in 1 after a finite number of coordinates starting at ω_{n+2} (after the first n + 1 coordinates which are always equal to ε_j for every element of Ω_{ε_j}). It is clear that this is a countable dense subset of Ω_{ε_j} . Let $C_{\varepsilon_j}^1$ consist of all the right end points that appear in the decreasing sequence of closed intervals $E_{\varepsilon_j}^k$ (k = 0, 1, 2, ...) whose intersection is the SVC set C_{ε_j} . Observe that $C_{\varepsilon_j}^1$ is also a countably infinite dense subset of C_{ε_j} .

We contend that all the right end points that comprise $C_{\varepsilon_j}^1$ are indexed by subscripts which are sequences in $\Omega_{\varepsilon_j}^1$. To see that this is so, observe that the first time that the terminal binary digit 1 appears as a coordinate in the subscript that indexes a given right end point in C_{ε_j} , in the *N*th coordinate say, counted after ε_j (which always appears

139

as the first n + 1 coordinates of every element in C_{ε_j}) so that $\omega_N = 1$ and $\omega_{N-1} = 0$ for N > n + 2, is in the set $E_{\varepsilon_j}^{N-n-1}$ at which the right end point $\underbrace{r_{\varepsilon_j,\ldots,\omega_{N-1},1}}_{N \text{ subscripts}}$ appears. Now

observe that after the infinite intersection $C_{\varepsilon_j} = \bigcap_{k=0}^{\infty} E_{\varepsilon_j}^k$ has been taken, this right end point $r_{\varepsilon_j,\ldots,\omega_{N-1},1}$ in C_{ε_j} is indexed by the subscript

N subscripts

$$r_{\varepsilon_j,\ldots,\omega_{N-1},1,1,1,\ldots}$$
, where $(\varepsilon_j,\ldots,\omega_{N-1},1,1,1,\ldots) \in \Omega^1_{\varepsilon_j}$

Thus, every right end point of each interval $E_{\varepsilon_i}^n$ is indexed by a subscript in $\Omega_{\varepsilon_i}^1$.

We define an order-preserving bijection $\xi_{\varepsilon_i}^1 : \Omega_{\varepsilon_i}^1 \to C_{\varepsilon_i}^1$ by

$$\xi_{\varepsilon_{j}}^{\mathbf{l}}(\varepsilon_{j},\omega_{n+2},\ldots,\omega_{N-1},1,1,1,1,\ldots)=r_{\varepsilon_{j},\omega_{n+2},\ldots,\omega_{N-1},111\ldots}$$

This map is plainly an order-preserving bijection, and extends to an order-preserving homeomorphism $\xi_{\varepsilon_i} : \Omega_{\varepsilon_i} \to C_{\varepsilon_i}$ as the set $\Omega^1_{\varepsilon_i}$ is dense in Ω_{ε_i} and $C^1_{\varepsilon_i}$ is dense in C_{ε_i} .

Now, consider the mapping $r : \Omega \to \Omega$ defined by $r((\omega_k)_{k\geq 1}) = (r(\omega_k)_{k\geq 1})$ where, for every $k \geq 1$,

$$r(\omega_k) = \begin{cases} 0 & \text{if } \omega_k = 1, \\ 1 & \text{if } \omega_k = 0. \end{cases}$$

It is immediate that r is an order-reversing homeomorphism and that

$$r(\Omega_{\varepsilon_i}) = \Omega_{\varepsilon_{(2^{n+1}+1)-i}}, \text{ for } 1 \le j \le 2^n.$$

To sum up, for all j with $2^n + 1 \le j \le 2^{n+1}$,

$$C_{\varepsilon_{j}} \xrightarrow{\xi_{\varepsilon_{j}}^{r-1}} \Omega_{\varepsilon_{j}} \xrightarrow{r} \Omega_{\varepsilon_{(2^{n+1}+1)-j}} \xrightarrow{Z_{\varepsilon_{(2^{n+1}+1)-j}}} Z_{\varepsilon_{(2^{n+1}+1)-j}}(\Omega_{\varepsilon_{(2^{n+1}+1)-j}}) \xrightarrow{\phi_{m}} K_{\varepsilon_{(2^{n+1}+1)-j}}$$

Since every mapping that appears in the sequence above is a homeomorphism, and all but one are order-preserving, the mapping $f_{\varepsilon_j,m,n} : C_{\varepsilon_j} \to K_{\varepsilon_{(2^{n+1}+1)-i}}$ defined by

$$f_{\varepsilon_j,m,n} = (\phi_m \circ Z_{\varepsilon_{(2^{n+1}+1)-j}} \circ r \circ \xi_{\varepsilon_j}^{-1}), \text{ for } 2^n + 1 \le j \le 2^{n+1}$$

must be an order-reversing homeomorphism.

To finish the construction of a continuous mapping in \mathcal{G}_m , define $\mathcal{S} = \bigcup_{j=2^{n+1}}^{2^{n+1}} C_{\varepsilon_j}$, $\mathcal{K} = \bigcup_{j=1}^{2^n} K_{\varepsilon_j}$ and $f_{m,n} : \mathcal{S} \to \mathcal{K}$ by setting $f_{m,n} \mid_{C_{\varepsilon_j}} = f_{\varepsilon_j,m,n}$. We define $f_{m,n}$ on $\mathcal{L}_m \cup \mathcal{R}_m$ to ensure that $f_{m,n} \in \mathcal{G}_m$ by setting

$$f_{m,n}(x) = \begin{cases} 3^m x & \text{for } x \in \mathcal{L}_m, \\ 3^m x - (3^m - 1) & \text{for } x \in \mathcal{R}_m. \end{cases}$$

We linearly interpolate to extend $f_{m,n}$ to \mathcal{M}_m from S. So, for any $s \in \mathcal{M}_m \setminus S$, we define $s_{\ell} = \sup(\{t \in S : t < s\} \cup \{1/3^m\}), s_r = \inf(\{t \in S : t > s\} \cup \{1 - 3^{-m}\})$ and

$$f_{m,n}(s) = f_{m,n}(s_{\ell}) + \frac{s - s_{\ell}}{s_r - s_{\ell}} (f_{m,n}(s_r) - f_{m,n}(s_{\ell})).$$

3.3. The proofs of the ancillary results. In this subsection, we furnish the proof of Proposition 3.1. To this end, we first state and prove two lemmas. These lemmas will then be used to establish Proposition 3.6, which itself will be used in the proof of Proposition 3.1.

LEMMA 3.4. For any positive integer m, let $\omega \in \Omega$ and $g_m \in \mathcal{G}_m$. Then, for any positive integer k, $g_m^k \circ \phi_m(\omega) = \phi_m(\sigma^k(\omega))$.

PROOF OF LEMMA 3.4. We argue by mathematical induction on k. If k = 1, then

$$g_m \circ \phi_m(\omega) = g_m \left(\sum_{k=1}^{\infty} \frac{(3^m - 1)\omega_k}{3^{mk}} \right) = 3^m \left(\sum_{k=1}^{\infty} \frac{(3^m - 1)\omega_k}{3^{mk}} \right) - (3^m - 1)\omega_1$$
$$= \sum_{k=2}^{\infty} \frac{(3^m - 1)\omega_k}{3^{m(k-1)}} = \sum_{k=1}^{\infty} \frac{(3^m - 1)\omega_{k+1}}{3^{mk}} = \phi_m(\sigma(\omega)).$$

Assume now, as our induction hypothesis, that for any positive integer $k \ge 1$,

$$g_m^k \circ \phi_m(\omega) = \phi_m(\sigma^k(\omega))$$

Then

140

$$g_m^{k+1} \circ \phi_m(\omega) = (g_m \circ g_m^k) \circ \phi_m(\omega) = g_m \circ (g_m^k \circ \phi_m)(\omega)$$
$$= g_m \circ \phi_m(\sigma^k(\omega)) = \phi_m(\sigma(\sigma^k(\omega))) = \phi_m(\sigma^{k+1}(\omega)).$$

Hence, the lemma follows by mathematical induction.

LEMMA 3.5. Let both $\omega = (\omega_k)_{k\geq 1}$ and $\tau = (\tau_k)_{k\geq 1}$ be elements of Ω . Then, for any positive integers m, k and N, the following three statements hold.

- (i) If $\omega_{k+j} = \tau_{k+j}$ for $1 \le j \le N$, then $|g_m^k(\phi_m(\omega)) g_m^k(\phi_m(\tau))| \le 3^{-mN}$. (ii) If $\omega_{k+j} = 0$ for $1 \le j \le N$, then $|g_m^k(\phi_m(\omega))| = g_m^k(\phi_m(\omega)) \le 3^{-mN}$.

(iii) If $\omega_{k+j} = 1$ for $1 \le j \le N$, then $|g_m^k(\phi_m(\omega))| = g_m^k(\phi_m(\omega)) \ge 1 - 3^{-mN}$.

PROOF OF LEMMA 3.5. From Lemma 3.4.

$$g_m^k(\phi_m(\omega)) = \phi_m(\sigma^k(\omega)) = \sum_{j=1}^{\infty} \frac{(3^m - 1)\omega_{k+j}}{3^{mj}}$$

and the three inequalities follow directly.

PROPOSITION 3.6. For any positive integers m and n, each of the 2^n mutually disjoint subsets K_{ε_i} (for $1 \le j \le 2^n$) is an extremally scrambled set for every function $g_m \in \mathcal{G}_m$.

PROOF OF PROPOSITION 3.6. In order to prove the proposition, we must show that each of the three defining conditions of an extremally scrambled set are satisfied by any particular K_{ε_i} for $1 \le j \le 2^n$ and for every $g_m \in \mathcal{G}_m$. To this end, choose any subset $\{x, y\} \subset K_{\varepsilon_i}$ such that x < y and any $g_m \in \mathcal{G}_m$. Assume also that x_0 is a periodic point of

 g_m having period p. Let $(\omega_x, \omega_y) \in Z_{\varepsilon_i}(\Omega_{\varepsilon_i}) \times Z_{\varepsilon_i}(\Omega_{\varepsilon_i})$ be such that

$$(\omega_x, \omega_y) = (\phi_m^{-1}(x), \phi_m^{-1}(y)).$$

Since ϕ_m is an order-preserving homeomorphism, ω_x and ω_y are unique elements of Ω , with $\omega_x \prec \omega_y$. Now since $\{\omega_x, \omega_y\} \subset Z_{\varepsilon_j}(\Omega_{\varepsilon_j})$ and $\omega_x \prec \omega_y$, to every positive integer u there corresponds a strictly increasing sequence $\{t_{u_r}\}_{r\geq 1}$ of positive integers such that $(\omega_x)_{t_{u_r}+j} = 0$ and $(\omega_y)_{t_{u_r}+j} = 1$ for $1 \le j \le u$. Thus, by Lemma 3.5,

$$\begin{split} |g_m^{t_{u_r}}(\phi_m(\omega_x)) - g_m^{t_{u_r}}(\phi_m(\omega_y))| &\geq |g_m^{t_{u_r}}(\phi_m(\omega_y))| - |g_m^{t_{u_r}}(\phi_m(\omega_x))| \\ &\geq \left(1 - \frac{1}{3^{mu}}\right) - \frac{1}{3^{mu}} = 1 - \frac{2}{3^{mu}}. \end{split}$$

Since

$$|g_m^{t_{u_r}}(\phi_m(\omega_x)) - g_m^{t_{u_r}}(\phi_m(\omega_y))| \ge 1 - \frac{2}{3^{mu}} \quad \text{for every } u \ge 1,$$

we conclude that

$$\lim_{k \to \infty} \sup_{k \to \infty} |g_m^k(x) - g_m^k(y)| = 1.$$

For the same pair $\{\omega_x, \omega_y\}$, since $\{\omega_x, \omega_y\} \subset Z_{\varepsilon_j}(\Omega_{\varepsilon_j})$, there is an increasing sequence $\{t_{u_s}\}_{s\geq 1}$ of positive integers such that $(\omega_x)_{t_{u_s}+j} = (\omega_y)_{t_{u_s}+j} = 0$ for $1 \leq j \leq u$. This implies, by Lemma 3.5, that

$$\begin{split} |g_m^{l_{u_s}}(\phi_m(\omega_x)) - g_m^{l_{u_s}}(\phi_m(\omega_y))| &\leq |g_m^{l_{u_s}}(\phi_m(\omega_x))| + |g_m^{l_{u_s}}(\phi_m(\omega_y))| \\ &\leq \frac{1}{3^{mu}} + \frac{1}{3^{mu}} = \frac{2}{3^{mu}}. \end{split}$$

As this estimate holds for every positive integer $u \ge 1$, we conclude that

$$\lim \inf_{k \to \infty} |g_m^k(x) - g_m^k(y)| = 0.$$

Finally, assume that x_0 is a periodic point of period p for the function g_m . We must prove that for any $x \in K_{\varepsilon_i}$,

$$\lim_{k \to \infty} \sup_{k \to \infty} |g_m^k(x) - g_m^k(x_0)| \ge \frac{1}{2}.$$

Observe now that $\max_{k\geq 1} |g_m^k(x_0)|$ is a definite real number in [0, 1] since we are simply choosing the maximum value from a finite set consisting of p distinct real numbers in [0, 1]. Assume that $\omega_x = \phi_m^{-1}(x) \in Z_{\varepsilon_j}(\Omega_{\varepsilon_j})$ where $x \in K_{\varepsilon_j}$. Then to every positive integer u there correspond strictly increasing sequences $(u_{r_i})_{t\geq 1}$ and $(u_{s_t})_{t\geq 1}$ of positive integers such that $(\omega_x)_{u_{r_i}+j} = 0$ and $(\omega_x)_{u_{s_t}+j} = 1$, for $1 \le j \le (u + p)$. Hence, by Lemma 3.5, $|g_m^{u_{r_t}}(x)| \le 3^{-m(u+p)}$ and

$$|g_m^{u_{r_t}+j}(x) - g_m^{u_{r_t}+j}(x_0)| \ge |g_m^{u_{r_t}+j}(x_0)| - |g_m^{u_{r_t}+j}(x)| \ge |g_m^{u_{r_t}+j}(x_0)| - \frac{1}{3^{mu}}$$

for $0 \le j \le p - 1$. Since the above inequalities hold for every positive integer *u*, we conclude that

$$\lim \sup_{k \to \infty} |g_m^k(x) - g_m^k(x_0)| \ge \max_{k \ge 1} |g_m^k(x_0)|$$

Similarly, by Lemma 3.5,

$$\begin{aligned} |g_m^{u_{s_t}}(x) - g_m^{u_{s_t}}(x_0)| &\ge |g_m^{u_{s_t}}(x)| - |g_m^{u_{s_t}}(x_0)| \\ &\ge \left(1 - \frac{1}{3^{mu}}\right) - |g_m^{u_{s_t}}(x_0)| \ge \left(1 - \frac{1}{3^{mu}}\right) - \max_{k \ge 1} |g_m^k(x_0)|.\end{aligned}$$

As this estimate holds for every positive integer *u*, we conclude that

$$\lim \sup_{k \to \infty} |g_m^k(x) - g_m^k(x_0)| \ge 1 - \max_{k \ge 1} |g_m^k(x_0)|.$$

Finally, since

$$\lim_{k \to \infty} \sup_{k \to \infty} |g_m^k(x) - g_m^k(x_0)| \ge \max_{k \ge 1} |g_m^k(x_0)| \quad \text{and}$$
$$\lim_{k \to \infty} \sup_{k \to \infty} |g_m^k(x) - g_m^k(x_0)| \ge 1 - \max_{k \ge 1} |g_m^k(x_0)|,$$

we have

$$\lim \sup_{k \to \infty} |g_m^k(x) - g_m^k(x_0)| \ge \frac{1}{2},$$

proving that K_{ε_j} must be an extremally scrambled set for any j with $1 \le j \le 2^n$ and for any particular $g_m \in \mathcal{G}_m$.

PROOF OF PROPOSITION 3.1. For any positive integers *n* and *k*, define n' = nk. From step 3 of the construction, we observe by definition that

$$f_{m,n'}(x) = \begin{cases} 3^m x & \text{for } x \in \mathcal{L}_m, \\ 3^m x - (3^m - 1) & \text{for } x \in \mathcal{R}_m. \end{cases}$$

Moreover, $f_{m,n'}$ is decreasing and continuous on both S and $\mathcal{M}_m^{\circ} \setminus S$ since it has been extended from S to \mathcal{M}_m by linear interpolation in such a way that $f_{m,n'}(1/3^m) = 1$ and $f_{m,n'}(1 - (1/3^m)) = 0$. Thus, $f_{m,n'}$ is in \mathcal{G}_m . Since $f_{m,n'} \in \mathcal{G}_m$, by Proposition 3.6, for any j with $2^{n'} + 1 \le j' \le 2^{n'+1}$, $K_{\varepsilon_{(2^{n'}+1_{+1})-j'}}$ must be an extremally scrambled set for $f_{m,n'}$, and since $f_{m,n'}(C_{\varepsilon_{j'}}) = K_{\varepsilon_{(2^{n'}+1_{+1})-j'}}$, for every $C_{\varepsilon_{j'}}$, it follows that every $C_{\varepsilon_{j'}}$ must also be an extremally scrambled set for $f_{m,n'}$. By construction it is clear that $\mu(C_{\varepsilon_{j'}}) = 1/2^{n'+2}$ and the $C_{\varepsilon_{j'}}$ are homomorphic to each other. Since n' = nk, for $1 \le j \le 2^n$, $M_{m,j}^n$ contains k mutually disjoint sets $M_{m,j'}^{n'}$ with $k(j-1) + 1 \le j' \le kj$, and thus k mutually disjoint extremally scrambled sets of identical positive Lebesgue measure, homeomorphic to each other. Thus, $f_{m,n'}$ is the desired map.

Acknowledgement

The authors are grateful to Minako Fujio and Aniruddha Ghosh for conversation and collaboration.

References

- F. Balibrea and V. Jiménéz-López, 'The measure of scrambled sets: a survey', *Acta Univ. M. Belii* Ser. Math. 7 (1999), 3–11.
- [2] F. Blanchard, W. Huang and L. Snoha, 'Topological size of scrambled sets', Colloq. Math. 110 (2008), 293–361.
- [3] L. Deng, M. Fujio and M. A. Khan, 'Eventual periodicity in the two-sector RSL model: equilibrium vis-à-vis optimum growth', *Econ. Theory* 72 (2021), 615–639.
- [4] O. Dovgoshey, O. Martio, V. Ryazanov and M. Vuorinen, 'The Cantor function', *Expo. Math.* 24 (2006), 1–37.
- [5] V. Jiménéz-López, 'Review of "Chaos on the Interval" by S. Ruette', *Mathematical Reviews* (2017), MR3616574.
- [6] I. Kan, 'A chaotic function possessing a scrambled set of positive Lebesgue measure', *Proc. Amer. Math. Soc.* 92 (1984), 45–49.
- [7] M. A. Khan and A. V. Rajan, 'On the eventual periodicity of piecewise linear chaotic maps', Bull. Aust. Math. Soc. 95 (2017), 467–475.
- [8] A. Y. Khinchin, *The Three Pearls of Number Theory* (Dover Publications, New York, 1998); translated from the second Russian edition published in 1948.
- [9] T. Li and J. A. Yorke, 'Period three implies chaos', Amer. Math. Monthly 82 (1975), 985–992.
- [10] J. Milnor and W. Thurston, 'On iterated maps of the interval', in: *Dynamical Systems*, J. C. Alexander (ed.), Lecture Notes in Mathematics, 1342 (Springer, Berlin, 1988), 465–563.
- M. Misiurewicz, 'One-dimensional dynamical systems', *Proc. Int. Cong. Math., Warszawa, 1983* Z. Ciesielski and C. Olech (eds.) (Polish Scientific Publishers PWN, Warsaw, 1984), 1277–1281.
- [12] M. Misiurewicz, 'Chaos almost everywhere', in: *Iteration Theory and its Functional Equations*, R. Liedl, L. Reich, and G. Targonski (eds.), Lecture Notes in Mathematics, 1163 (Springer, Berlin, 1985), 125–130.
- [13] M. B. Nathanson, 'Piecewise linear functions with almost all points eventually periodic', Proc. Amer. Math. Soc. 92 (1976), 75–81.
- [14] Y. Oono, *The Nonlinear World: Conceptual Analysis and Phenomenology*, Springer Series in Synergetics (Springer, Tokyo, 2013).
- [15] A. V. Rajan, 'Positive measure scrambled sets of some chaotic functions', *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* 28 (2018), Article no. 1850052.
- [16] S. Ruette, *Chaos on the Interval*, University Lecture Series, 67 (American Mathematical Society, Providence, RI, 2017); reviewed by V. Jiménéz-López.
- [17] O. M. Sharkovsky, 'Co-existence of cycles of a continuous mapping of the line into itself', Ukrainian Math. J. 16 (1964), 61–71.
- [18] J. Smítal, 'A chaotic function with a scrambled set of positive Lebesgue measure', Proc. Amer. Math. Soc. 92 (1984), 50–54.

LIUCHUN DENG, Social Sciences Division, Yale-NUS College, Singapore e-mail: yncdl@nus.edu.sg

M. ALI KHAN, Department of Economics, The Johns Hopkins University, Baltimore, MD 21218, USA e-mail: akhan@jhu.edu

ASHVIN V. RAJAN, 3935 Cloverhill Road, Baltimore, MD 21218, USA e-mail: ashvinrj@aol.com